Gröbner Bases Native to 'pseudo'-Hodge Algebras, with Application to the Algebra of Minors

Abhiram Natarajan

Joint work with Joshua Grochow (Univ. of Colorado, Boulder, USA)

▶ In Computational Complexity Theory, you usually have sets parameterized by $n \in \mathbb{Z}$, and seek asymptotic information that is symbolic in n, e.g. complexity of $n \times n$ matrix multiplication

- ▶ In Computational Complexity Theory, you usually have sets parameterized by $n \in \mathbb{Z}$, and seek asymptotic information that is symbolic in n, e.g. complexity of $n \times n$ matrix multiplication
- Sometimes you get intuition by computing small examples, e.g. n = 2, 3, ...

- ▶ In Computational Complexity Theory, you usually have sets parameterized by $n \in \mathbb{Z}$, and seek asymptotic information that is symbolic in n, e.g. complexity of $n \times n$ matrix multiplication
- Sometimes you get intuition by computing small examples, e.g. n = 2, 3, ...
- ► Gröbner bases are well-suited to both of the above!

- ▶ In Computational Complexity Theory, you usually have sets parameterized by $n \in \mathbb{Z}$, and seek asymptotic information that is symbolic in n, e.g. complexity of $n \times n$ matrix multiplication
- Sometimes you get intuition by computing small examples, e.g. n = 2, 3, ...
- Gröbner bases are well-suited to both of the above!
- ► Gröbner bases give theoretical insight as well as are the key tool in effective methods

Gröbner Bases are not a Panacea

 Obtaining a Gröbner basis can be tremendously expensive -EXPSPACE-complete (Mayr and Meyer [1982])

Gröbner Bases are not a Panacea

▶ Obtaining a Gröbner basis can be tremendously expensive -EXPSPACE-complete (Mayr and Meyer [1982])

► Hard to get even simple cases to finish, e.g. 3×3 determinant orbit closure, tensor rank of 3×3 multiplication

Gröbner Bases are not a Panacea

▶ Obtaining a Gröbner basis can be tremendously expensive -EXPSPACE-complete (Mayr and Meyer [1982])

▶ Hard to get even simple cases to finish, e.g. 3×3 determinant orbit closure, tensor rank of 3×3 multiplication

Gröbner bases tend to obscure symmetry!

► There are Gröbner basis methods to obtain Algebraic de Rham Cohomology (Oaku and Takayama [1999, 2001])

- ► There are Gröbner basis methods to obtain Algebraic de Rham Cohomology (Oaku and Takayama [1999, 2001])
- ► Requires computing Gröbner bases of D-ideals (ideals in the Weyl algebra):

$$W_{n} := \mathbb{C}\left[\left\{X_{i}, \frac{\partial}{\partial X_{i}}\right\}_{i \in [n]}\right] \left/\left\langle\left\{\frac{\partial}{\partial X_{i}} \cdot X_{i} - X_{i} \cdot \frac{\partial}{\partial X_{i}} - 1\right\}_{i \in [n]}\right\rangle$$

- ► There are Gröbner basis methods to obtain Algebraic de Rham Cohomology (Oaku and Takayama [1999, 2001])
- ► Requires computing Gröbner bases of D-ideals (ideals in the Weyl algebra):

$$W_n := \mathbb{C}\left[\left\{X_i, \frac{\partial}{\partial X_i}\right\}_{i \in [n]}\right] \left/ \left\langle \left\{\frac{\partial}{\partial X_i} \cdot X_i - X_i \cdot \frac{\partial}{\partial X_i} - 1\right\}_{i \in [n]} \right\rangle$$

We couldn't even obtain cohomology of 3 x 3 determinant hypersurface by hand, or on Macaulay2

- ► There are Gröbner basis methods to obtain Algebraic de Rham Cohomology (Oaku and Takayama [1999, 2001])
- ► Requires computing Gröbner bases of D-ideals (ideals in the Weyl algebra):

$$W_{n} := \mathbb{C}\left[\left\{X_{i}, \frac{\partial}{\partial X_{i}}\right\}_{i \in [n]}\right] \left/\left\langle\left\{\frac{\partial}{\partial X_{i}} \cdot X_{i} - X_{i} \cdot \frac{\partial}{\partial X_{i}} - 1\right\}_{i \in [n]}\right\rangle$$

► We couldn't even obtain cohomology of 3 × 3 determinant hypersurface by hand, or on Macaulay2

Question

Develop a Gröbner basis theory which takes advantage if variety corresponding to ideal has large symmetry group, or is 'determinantal'

Hodge Algebra (Alg. with Straightening Law)

- ▶ If an algebra A is a pseudo-ASL (p-ASL for short) then:
 - $ightharpoonup A \cong \mathbb{F}[\vec{X}] / I$, and

Flodge Algebra (Alg. with Straightening Law)

- ▶ If an algebra A is a pseudo-ASL (p-ASL for short) then:
 - $ightharpoonup A \cong \mathbb{F}[\vec{X}] / I$, and
 - ▶ there is a monomial ideal $\Sigma \subseteq \mathbb{F}[\vec{X}]$ such that monomials not in Σ form an \mathbb{F} -linear basis for A; such monomials are called standard monomials.

Hodge Algebra (Alg. with Straightening Law)

- ▶ If an algebra A is a pseudo-ASL (p-ASL for short) then:
 - $ightharpoonup A \cong \mathbb{F}[\vec{X}] / J$, and
 - ▶ there is a monomial ideal $\Sigma \subseteq \mathbb{F}[\vec{X}]$ such that monomials not in Σ form an \mathbb{F} -linear basis for A; such monomials are called standard monomials.
- ► If A is an ASL, the product of two standard monomials can be straightened into a linear combination of 'smaller' standard monomials

Hodge Algebra (Alg. with Straightening Law)

- ▶ If an algebra A is a pseudo-ASL (p-ASL for short) then:
 - $ightharpoonup A\cong \mathbb{F}[\vec{X}]\Big/J$, and
 - ▶ there is a monomial ideal $\Sigma \subseteq \mathbb{F}[\vec{X}]$ such that monomials not in Σ form an \mathbb{F} -linear basis for A; such monomials are called standard monomials.
- ► If A is an ASL, the product of two standard monomials can be straightened into a linear combination of 'smaller' standard monomials
- ► ASLs arise as coordinate rings of algebraic varieties, e.g. Grassmanians, determinantal varieties, flag varieties, Schubert varieties

Bideterminants (products of minors)

► Example of Hodge algebra - algebra of bideterminants

e.g.
$$A = \mathbb{F}[X_{1,1}, X_{1,2}, X_{2,1}, X_{2,2}, Y] / \langle X_{1,2} X_{2,1} - X_{1,1} X_{2,2} + Y \rangle$$

$$\Sigma = \langle X_{1,2}X_{2,1}\rangle\,; \qquad \text{(turns out } A \cong \mathbb{F}[X_{1,1},X_{1,2},X_{2,1},X_{2,2}])$$

Bideterminants (products of minors)

► Example of Hodge algebra - algebra of bideterminants

$$\begin{split} \text{e.g.} \quad & A = \mathbb{F}[X_{1,1}, X_{1,2}, X_{2,1}, X_{2,2}, Y] \left/ \left\langle X_{1,2} X_{2,1} - X_{1,1} X_{2,2} + Y \right\rangle \right. \\ & \Sigma = \left\langle X_{1,2} X_{2,1} \right\rangle; \qquad \text{(turns out } A \cong \mathbb{F}[X_{1,1}, X_{1,2}, X_{2,1}, X_{2,2}]) \end{split}$$

- ► The above generalizes -
 - \triangleright poly ring with one variable for each minor of $n \times m$ matrix
 - quotient by relations between minors
 - \blacktriangleright gives ASL structure to the co-ordinate ring of $n \times m$ matrices
 - > standard monomials correspond to standard bitableaux

Bideterminants (products of minors)

Example of Hodge algebra - algebra of bideterminants

$$\begin{split} \text{e.g.} \quad & A = \mathbb{F}[X_{1,1}, X_{1,2}, X_{2,1}, X_{2,2}, Y] \left/ \left\langle X_{1,2} X_{2,1} - X_{1,1} X_{2,2} + Y \right\rangle \right. \\ & \quad & \Sigma = \left\langle X_{1,2} X_{2,1} \right\rangle; \qquad \text{(turns out } A \cong \mathbb{F}[X_{1,1}, X_{1,2}, X_{2,1}, X_{2,2}]) \end{split}$$

- ► The above generalizes
 - ightharpoonup poly ring with one variable for each minor of $n \times m$ matrix
 - quotient by relations between minors
 - \blacktriangleright gives ASL structure to the co-ordinate ring of $\mathfrak{n} \times \mathfrak{m}$ matrices
 - ► standard monomials correspond to standard bitableaux
- ► Advantage smaller expressions for 'determinant-like' polynomials; bideterminants are reflect symmetries coming from the action (representation theory) of GL_n

► Generally, in p-ASLs, you have smaller expressions for polynomials that are adapted to the p-ASL basis

- Generally, in p-ASLs, you have smaller expressions for polynomials that are adapted to the p-ASL basis
- ► Can we have a theory of Gröbner bases directly in p-ASLs?

- ► Generally, in p-ASLs, you have smaller expressions for polynomials that are adapted to the p-ASL basis
- ► Can we have a theory of Gröbner bases directly in p-ASLs?
- Trivially, any ASL $A\cong \mathbb{F}[\vec{X}] \Big/J$, so for an any ideal $I\subseteq A$, we can look Gröbner bases of the ideal $I+J\subseteq \mathbb{F}[\vec{X}]$ this obscures symmetries

- ► Generally, in p-ASLs, you have smaller expressions for polynomials that are adapted to the p-ASL basis
- Can we have a theory of Gröbner bases directly in p-ASLs?
- Trivially, any ASL $A\cong \mathbb{F}[\vec{X}] \Big/J$, so for an any ideal $I\subseteq A$, we can look Gröbner bases of the ideal $I+J\subseteq \mathbb{F}[\vec{X}]$ this obscures symmetries

Question

Can we build a theory of Gröbner bases 'native' to p-ASLs, i.e. Gröbner theory without referencing the ideal J?

Challenges

lacksquare Basis of p-ASL $A={\mathbb F}[ec X]\Big/J$ consists only of standard monomials (monomials outside Σ), not all monomials in ec X

Challenges

- lackbrack Basis of p-ASL $A=\mathbb{F}[\vec{X}]\Big/J$ consists only of standard monomials (monomials outside Σ), not all monomials in \vec{X}
- Product of standard monomials not necessarily standard, might require straightening

Challenges

- lackbrack Basis of p-ASL $A=\mathbb{F}^{[ec{X}]}\Big/J$ consists only of standard monomials (monomials outside Σ), not all monomials in $ec{X}$
- Product of standard monomials not necessarily standard, might require straightening
 - ► How do you define term order?
 - How would you define division of monomials?
 - ► What plays the role of monomial ideals?

Term Order & Division

- ightharpoonup A p-ASL term order on a p-ASL A is a total order \prec on standard monomials in A such that
 - ▶ 1 ≤ m
 - ▶ If $a \prec b$ and $c \leq d$, and ac, $bd \neq 0$, then

 $LM(ac) \prec LM(bd)$

Term Order & Division

- ightharpoonup A p-ASL term order on a p-ASL A is a total order \prec on standard monomials in A such that
 - ▶ 1 ≤ m
 - ▶ If $a \prec b$ and $c \preceq d$, and ac, $bd \neq 0$, then

$$LM(ac) \prec LM(bd)$$

- ► When does standard monomial m divide m':
 - ordinary division in the polynomial ring, or
 - ▶ m divides m' if there exists standard monomial f such that

$$LM(mf) = m'$$

Auxilliary Algebra of Leading Terms

► Given p-ASL A, algebra of leading terms w.r.t. A is another p-ASL A_{1t} on the same variables, and the same standard monomials such that for standard monomials m, m' no straightening

$$\pi_{lt}(\mathfrak{m})\cdot\pi_{lt}(\mathfrak{m}')=$$
 or $\underline{\pi_{lt}(LT(\mathfrak{mm}'))}$ leading term of straightening

where $\pi_{lt}:A \to A_{lt}$ is the identity

Auxilliary Algebra of Leading Terms

► Given p-ASL A, algebra of leading terms w.r.t. A is another p-ASL A_{1t} on the same variables, and the same standard monomials such that for standard monomials m, m' no straightening

$$\pi_{lt}(\mathfrak{m})\cdot\pi_{lt}(\mathfrak{m}')=$$
 or $\underline{\pi_{lt}(LT(\mathfrak{m}\mathfrak{m}'))}$ leading term of straightening

where $\pi_{lt}:A \to A_{lt}$ is the identity

Proposition

Every p-ASL A admits two algebras of leading terms – A_{gen} where the product is never 0, and, A_{disc} where product is 0 unless mm' is also a standard monomial.

Definition of p-ASL Gröbner Basis

- ▶ Given p-ASL A, algebra of leading terms A_{lt} , and an ideal $I \subseteq A$, then $G \subseteq A$ is a p-ASL Gröbner basis if:
 - ▶ For all $f \in I$, there exists $g \in G$ such that $\pi_{lt}(LM(g))$ divides $\pi_{lt}(LM(f))$, or

Definition of p-ASL Gröbner Basis

- ▶ Given p-ASL A, algebra of leading terms A_{lt} , and an ideal $I \subseteq A$, then $G \subseteq A$ is a p-ASL Gröbner basis if:
 - ▶ For all $f \in I$, there exists $g \in G$ such that $\pi_{lt}(LM(g))$ divides $\pi_{lt}(LM(f))$, or

Our Main Result

Theorem (Grochow-N, 2025)

For any p-ASLs A with a p-ASL term order, we have a theory of Gröbner bases native to A. Specifically:

Our Main Result

Theorem (Grochow-N, 2025)

For any p-ASLs A with a p-ASL term order, we have a theory of Gröbner bases native to A. Specifically:

	ordinary Gröbner	p-ASL Gröbner theory
Existence	│	√
Reduced	1	✓
Universal	1	1
Syzygies	1	o√ o√
Algorithms	1	
Krull Dim.	1	√

Our Main Result

Theorem (Grochow-N, 2025)

For any p-ASLs A with a p-ASL term order, we have a theory of Gröbner bases native to A. Specifically:

	ordinary Gröbner	p-ASL Gröbner theory
Existence	√	
Reduced	√	
Universal	✓	
Syzygies	✓	
Algorithms	✓	
Krull Dim.	√	✓

Corollary (Grochow-N, 2025)

The algebra of bideterminants has a p-ASL term order, thus we have a Gröbner basis theory (called bd-Gröbner bases).

Applications to Bideterminant Algebra

► Universal p-ASL Gröbner basis is a p-ASL Gröbner basis for any p-ASL term order and any algebra of leading terms

Applications to Bideterminant Algebra

► Universal p-ASL Gröbner basis is a p-ASL Gröbner basis for any p-ASL term order and any algebra of leading terms

Theorem (Grochow-N, 2025)

For any \mathfrak{r} , the set of minors of size $\geqslant \mathfrak{r}$ is a universal bd-Gröbner basis for its ideal.

Applications to Bideterminant Algebra

► Universal p-ASL Gröbner basis is a p-ASL Gröbner basis for any p-ASL term order and any algebra of leading terms

Theorem (Grochow-N, 2025)

For any \mathfrak{r} , the set of minors of size $\geqslant \mathfrak{r}$ is a universal bd-Gröbner basis for its ideal.

Takeaway

- 1. Given all our machinery, the proof is one-line
- 2. In the ordinary case, universal Gröbner basis are known only for maximal minors and minors of size 2

▶ In upcoming work, we have already extended our Gröbner basis theory to mildly non-commutative algebras, including the Weyl algebra

- ► In upcoming work, we have already extended our Gröbner basis theory to mildly non-commutative algebras, including the Weyl algebra
- ► Get bd-Gröbner bases of annihilating D-ideals

- In upcoming work, we have already extended our Gröbner basis theory to mildly non-commutative algebras, including the Weyl algebra
- ► Get bd-Gröbner bases of annihilating D-ideals
- ► Compute Weyl closure, b-functions, etc. using bd-Gröbner bases in the Weyl algebra

- In upcoming work, we have already extended our Gröbner basis theory to mildly non-commutative algebras, including the Weyl algebra
- ► Get bd-Gröbner bases of annihilating D-ideals
- ► Compute Weyl closure, b-functions, etc. using bd-Gröbner bases in the Weyl algebra
- ➤ See if we can develop a bipermanent Gröbner basis theory (codimension of singular locus of permanent hypersurface is unknown!)

References

- E. W. Mayr and A. R. Meyer. The complexity of the word problems for commutative semigroups and polynomial ideals. Advances in mathematics, 46(3):305-329, 1982.
- T. Oaku and N. Takayama. An algorithm for de Rham cohomology groups of the complement of an affine variety via D-module computation. J. Pure Appl. Algebra, 139(1-3):201-233, 1999. doi: 10.1016/S0022-4049(99)00012-2.
- T. Oaku and N. Takayama. Computing de Rham cohomology groups. In Proceedings of the 33rd Symposium on Ring Theory and Representation Theory (Shimane, 2000), pages 19-22. Tokyo Univ. Agric. Technol., Tokyo, 2001.